arXiv:1210.3100v1 [cond-mat.mes-hall] 11 Oct 2012

Localized and polarized charged zero modes in three-dimensional topological insulators induced by a magnetic vortex

J.M. Fonseca[*]
Grupo de Física/ICET/CRP, Universidade Federal de Viçosa,
Rodovia MG-230 Km 7, Cep 38810-000, Rio Paranaíba, Minas Gerais, Brazil.

W.A. Moura-Melo[†] and A.R. Pereira[‡]
Departamento de Física, Universidade Federal de Viçosa,
Viçosa, 36570-000, Minas Gerais, Brazil

(Dated: October 12, 2012)

Coating a three-dimensional topological insulator with a ferromagnetic film supporting an in-plane magnetic vortex renders electronic states bound to the vortex. Charged zero-modes also appear as fully polarized current localized near the vortex center. From the magnetic point of view, such a special current can be faced as an alternative technique for detecting magnetic vortices in thin films.

PACS numbers: 73.20.-r, 75.30.Hx, 85.75.-d

I. INTRODUCTION

Topological Insulator (TI) is a nontrivial class of band insulator characterized by a topological index associated with the Bloch wave-function in the momentum space. They are materials with a bulk band gap generated by strong spin-orbit coupling and topologically protected metallic surface states. At present, such a behavior has been observed in Bi_xSb_{1-x} alloys, Bi_2Se_3, Bi_2Te_3, Sb_2Se_3 crystals[1–3] and in ternary intermetallic Heusler compounds[4,5]. The reason for the considerable interest in topological insulators lies in their potential for producing new physical phenomena as well as future technological applications. These rely on the special properties of the topologically protected electronic surface states, composed of an odd number of Dirac fermions, familiar to the physics of graphene[6]. The physical manifestation of the topological order comes in the form of protected, gapless surface states that are robust against damaging the surface by chemical or mechanical means such as alterations in its shape or orientation with respect to the crystal lattice, or even by disordering the bulk, as long as such changes are applied in moderation[7]. The above statements apply exclusively to systems that respect time-reversal-symmetry (TRS), that is, nonmagnetic TI in zero external magnetic field. When TRS is broken, even by a weak magnetic perturbation, a gap can be open up into the spectrum of the topologically protected surface states. When a gap opens as a consequence of a magnetic perturbation, the resulting surface is not an ordinary insulator; instead, it is a quantum Hall insulator with properties similar to those of the familiar quantum Hall systems realized in two-dimensional electron gas[8]. A tunable energy gap at the surface Dirac point provides a means to control the surface electric transport, which is of great importance for applications.

Topological order in proximity to magnetism has been considered as one of the main interest in the field[8–13]. In principle, it can be realized by several means such as the application of weak magnetic fields through a Zeeman term[14] $H_Z = -g\mu_B \vec{s}\cdot\vec{B}$, or through exchange coupling to magnetic thin films[11] $H_{int} = -\Delta\vec{M}\cdot\vec{s}$, and magnetic impurities[8,15,16] $H_{imp} = \sum_i \vec{S}_i\cdot\vec{s}\delta(\vec{r}-\vec{R}_i)$. The exchange coupling arises due to the spin-spin interaction between spins in different atoms. Such an interaction is due to the superposition (overlap) between the wavefunction that describes conduction electrons on the surface of a TI and bound electrons in the magnetic impurity on the surface. This is one of the most appealing from the theoretical point of view because it not only gives a simple mechanism to develop the theory of the topologically quantized magnetoelectric term, but it also allows us to look for unexpected effects that can alter the magnetization dynamics[12,13]. Therefore, here we consider a 3D TI coated by thin film of layered two-dimensional ($2D$) Heisenberg ferromagnet. Indeed, magnetic materials (ferromagnetic or even antiferromagnetic) in two spatial dimensions may support topological excitations such as skyrmions and vortices. Particularly, vortices arise in classical magnetic systems containing an easy-plane anisotropy, which makes the spins to prefer to point along the XY-plane. For instance, easy-plane ferromagnets are described by the Hamiltonian $H = -J\sum_{i,j}[S_i^x S_j^x + S_i^y S_j^y + \lambda S_i^z S_j^z]$, where $J > 0$ is the exchange constant, $0 \leq \lambda < 1$ is the easy-plane anisotropy and $\vec{S}_i = (S_i^x, S_i^y, S_i^z)$ is the classical spin vector at site i. The spin field can be parametrized by two scalar fields ϕ and $\varepsilon = \cos\theta$ (the azimuthal and polar angles) as follows: $\vec{S} = \{\sqrt{1-\varepsilon^2}\cos\phi, \sqrt{1-\varepsilon^2}\sin\phi, \varepsilon\}$. Taken into account the most realistic cases of discrete lattices, and depending on the range of λ, such an easy-plane system supports two types of static vortices: the in-plane vortex (in which all spins are confined to the XY-plane[17]) and the out-of-plane vortex (in which some spins around the vortex center can point perpendicularly to the XY-plane). Indeed, considering a critical value of

λ denoted by λ_c, then, for the range $\lambda < \lambda_c$, the stable excitation is the in-plane vortex while, for $\lambda > \lambda_c$, the out-of-plane vortex becomes stable[18]. The stability of these solutions has only been determined via computer simulations. The critical anisotropy λ_c depends on the lattice geometry: for the square lattice, $\lambda_c = 0.72$; similarly $\lambda_c = 0.86$ for the hexagonal lattice and $\lambda_c = 0.62$ for the triangular lattice[18]. Qualitatively, similar results can be obtained for $2D$ easy-plane antiferromagnetic systems.

Our interest here is to investigate how an in-plane ferromagnetic vortex affects the electronic states lying on the surface of a 3D TI. Although $2D$ magnetic materials can support vortex excitations, there is, however, the experimental challenging of getting and confining a single static vortex in these thin films[19–21]. Even the observation of these excitations in $2D$ layered easy-plane magnetic materials is not a simple task: Indeed, only indirect evidences for their appearance has been reported so far[19,22,23]. In addition, there is the problem of getting a static single vortex living in the system for a long time[24]. To overcome these difficulties, a possibility should be the use of a vortex pinning mechanism. In $2D$ magnetic materials, a nonmagnetic impurity (a vacancy) is able to pin the vortex center[25–29], localizing the excitation around it[25,27,28,30]. Another possibility (perhaps the simplest) should be through the use of thin magnetic nanodisks over the TI. Really, depending on the size of a nanodisk, magnetic vortices are naturally the ground state of these systems and they can be directly observed by experimental techniques[31,32]. Moreover, the vortex ground state is always out-of-plane (in the vortex core, the spins point perpendicularly to the disk plane) due to the competition between exchange and dipolar interactions. However, the presence of artificial defects intentionally inserted in the nanodisk (such as holes)[33–36], may destroy the usual out-of plane spin structure in the vortex core (after vortex capture), resulting in an essentially confined planar vortex centered at the hole[37].

Here we are interested in the in-plane vortex effects on the TI. Then, in principle, it can be investigated either by applying a thin coating of magnetic material (with $\lambda < \lambda_c$ and containing a small percentage of nonmagnetic impurities) on the TI or by putting a magnetic nanodisk (with a hole, preferentially in the disk center) on the top of the TI. In the context of the first situation, recently, the authors of Ref.[13] have theoretically studied the magnetic textures, such as domain walls and vortices in a ferromagnetic thin film deposited on the surface of a TI. These authors have found that the magnetic textures interacting with the topological surface states are electrically charged. The experimental observation of massive Dirac Fermion on the surface of a magnetically doped TI was reported in Ref.[8], where the authors introduced magnetic dopants into three-dimensional TI Bi_2Se_3 and observed the massive Dirac fermion state by angle-resolved photoemission. Other aspects as Hedgehog spin texture[38], magnetic scattering of the carriers on the surface[39], magnetic ordering[40,41] and topological magneto-electric effect[10,11,42] were also investigated.

Considering an in-plane closed flux magnetized state interacting with a 3D TI, we have realized that the interaction between the surface carriers and the vortex leads to bound-states and spin polarized zero-modes, besides scattered states. Such a framework provides a mechanism to obtain spin polarized current based on the zero-mode states; conversely, such a physical mechanism appears to be useful for experimental magnetism to precise and efficiently detection of magnetic vortices in layered magnetic thin films. Our article is outline as follows: in Section II we present the model which takes in to account the basic properties of the TI surface electronic states along with their coupling to the magnetic moments (spins) lying on the ferromagnetic layer. Section III is devoted to the zero-mode states induced by the vortex. Besides its basic features we also point out possible utilities for such a special current. In Section IV the remaining bound-states are presented and discussed, completing the 'atomic picture' provided by the compound system vortex-TI charge carries. We finally close our paper by pointing out our conclusions and prospects for future investigation.

II. THE MODEL

The low energy excitations on the surface of a TI are described by a Dirac-like equation in a two-dimensional plane. When the chemical potential is confined between valence and conduction bands (in the ground state), the positive energy modes (that describe free conduction electrons) are empty, while the negative energy modes (that describe bound valence electrons) are filled with valence electrons. The mass term in the equation expresses the gap of energy between the conduction and valences bands (a trivial insulator) and a vanishing mass term leads to a gapless spectrum conductor or TI surface modes[43]. The latter ones offer a unique platform to investigate the physics of robust Dirac points.

Here, we analyze the simplest case of a single Dirac point (for instance, Bi_2Se_3, Bi_2Te_3 and Sb_2Te_3[44]). Let us consider the surface of the $3D$ TI in the XY-plane (its bulk extends to $z < 0$ region) coated by a ferromagnetic thin film supporting an in-plane vortex. Therefore, the low-energy effective Hamiltonian for the TI surface states coupled to the vortex magnetic moments (spins) reads:

$$H = H_0 + H_{int} \tag{1}$$

where

$$H_0 = v_F \psi^\dagger(\vec{r})(k_x\sigma^x + k_y\sigma^y)\psi(\vec{r})\,, \tag{2}$$

and

$$H_{int} = -\sum_{i=x,y,z} \Delta_i \vec{M}_i(\vec{r}) \cdot \vec{s}(\vec{r})\,. \tag{3}$$

Above, $\vec{s}(\vec{r}) = \psi^\dagger(\vec{r})\vec{\sigma}\psi(\vec{r})$ is the spin density of surface electrons in the position $\vec{r}$, $\vec{M}(\vec{r})$ is the magnetic moment applied on the surface of the TI, which characterizes the magnetic configuration on the surface and Δ_i is the ferromagnetic exchange constant[7,9], accounting for the coupling between the electrons lying on the TI surface and the magnetic moments composing the magnetic material (exchange-type proximity effect; note that H_{int} is minimized when the electrons spins align to the dipoles). For instance, regarding the material Sb_2Te_3, doped with vanadium the surface exchange parameters Δ_x, Δ_y, Δ_z are estimated to be of the order of $0.1-0.5eV$, relying on the overlap between the wave-functions[9]. For the case of a ferromagnetic ordered state perpendicular to the surface, the interaction Hamiltonian (3) has the form of a mass term for the surface electrons proportional to σ_z, being the carriers massive with a dispersion relation like:

$$E = \pm v_F\sqrt{k_x^2 + k_y^2 + (\mu v_F)^2}\,, \tag{4}$$

where the positive and negative signs describe the conduction electrons and the valence electrons (or holes), respectively. Whenever this is in order a gap opens, $E_g = 2\mu v_f^2$, rendering TI to be a quantum Hall insulator. As we shall show below, the magnetic vortex also breaks TRS in the surface, but does not induces a mass gap, like above. As consequences we shall find electronic bound-states and zero-energy modes around the vortex.

A. Magnetic vortex configuration

An in-plane vortex in the continuum limit may be described by:

$$\varepsilon = \varepsilon_v = 0, \quad \phi = \phi_v = q\arctan\left[\frac{y-y_0}{x-x_0}\right] \pm \phi_0, \tag{5}$$

where q accounts for its topological charge (vorticity), (x_0, y_0) is its centered position while ϕ_0 is a constant determining its profile (in Fig.1 we depict two examples of unity charge vortex, $q = 1$, with distinct profiles). For such a planar vortex, its magnetization, $\vec{M}$, is essentially (in polar coordinates): $\vec{M}(\vec{r}) = M\hat{\phi}$ with $M^2 = M_x^2 + M_y^2 =$ constant and therefore, the interaction Hamiltonian (3) becomes:

$$H_{int} = -\psi^\dagger(\vec{r})M\hat{\phi}\cdot\vec{\sigma}\psi(\vec{r})\,, \tag{6}$$

with $M = \Delta_i M_i$ (hereafter we normalize $\Delta_i \equiv 1\forall i$). This term does not open a mass gap in the spectrum, like an ordered ferromagnet does; It rather induces the appearance of bound-states, including zero-modes, in the electronic spectrum of the composed system, as we shall see below. Thus, the dynamics of the low energy states is described, in polar coordinates, by the Hamiltonian:

$$\mathcal{H} = -i\hbar v_F \times \begin{pmatrix} 0 & e^{-i\theta}\left(\frac{\partial}{\partial r} - \frac{i}{r}\frac{\partial}{\partial\phi} - \frac{M}{\hbar v_F}\right) \\ e^{i\theta}\left(\frac{\partial}{\partial r} + \frac{i}{r}\frac{\partial}{\partial\phi} + \frac{M}{\hbar v_F}\right) & 0 \end{pmatrix} \tag{7}$$



$\phi_0 = \pi/2$ $\phi_0 = 0$

FIG. 1. (Color online) Examples of ferromagnetic vortex-like patterns with unity charge, $q = +1$, and with profiles $\phi_0 = +\pi/2$ (curly-type) and $\phi_0 = 0$ (radial-like). The depicted arrows represent the magnetization vector field, $\vec{M}$.

Thus, the eigenvalue problem $\mathcal{H}\psi = E\psi$ can be solved analytically by the usual ansatz:

$$\psi(\vec{r}) = \begin{pmatrix} e^{im\phi}\varphi(r) \\ e^{i(m+1)\phi}\chi(r) \end{pmatrix}\,, \quad m = 0, \pm 1, \pm 2, \ldots\,, \tag{8}$$

where the components of the spinor $\varphi(r)$ and $\chi(r)$ satisfy the first order coupled differential equations given by:

$$\left(\frac{d}{dr} + \frac{m+1}{r} - \frac{M}{\hbar v_F}\right)\chi(r) = i\frac{E}{\hbar v_F}\varphi(r)\,, \tag{9}$$

$$\left(\frac{d}{dr} - \frac{m}{r} + \frac{M}{\hbar v_F}\right)\varphi(r) = i\frac{E}{\hbar v_F}\chi(r)\,, \tag{10}$$

Such expressions may be reduced to a second order decoupled differential equation, but to analyze zero modes solutions, this form is more appropriated, as below.

III. ZERO-ENERGY MODES

Firstly, we investigate the zero energy solutions (zero-modes) of the equations (9) and (10) for the case of a planar vortex with $\phi_0 = \pi/2$. When the Dirac equation has a homogeneous position independent mass term, the solutions are as usual: continuum solutions with positive ($E > 0$) and negative ($E < 0$) energies. However, in the presence of a topologically non-trivial background or defect, a mass term position dependent can appear and the spectrum contains (in addition to the habitual positive and negative energy modes) also isolated zero-energy mode states. The zero-energy solutions of the Dirac equation do not require a specific form for the inhomogeneous mass profile; instead, they only belong to a nontrivial topological class. Then, the presence of zero-energy modes can be established, a priori, by mathematical index theorems, which relate the occurrence of these zero modes to the geometry and topology of the space on which the Dirac equation is stated[43]. The index theorem offers an analytic tool that relates the zero modes of elliptic operators (like the Dirac Hamiltonian in (2+1) dimensions) with the geometry of the manifold on which these operators are defined[45]. This theorem can be employed to obtain information about the low energy behavior of the TI and, in particular, about its conductivity properties. A general study of the index theorem in TI will be presented elsewhere[46].

Now, we investigate the possibility of occurrence of zero-modes on a surface of the TI without a mass profile in the Dirac equation due to the presence of a non-trivial background for the surface carriers. We study this occurrence based on analytical procedures by explicitly solving the Dirac equation. In the case of zero energy solutions, the radial components of the spinors (9) and (10) must satisfy:

$$\left(\frac{d}{dr} - \frac{m}{r} + \frac{M}{\hbar v_F}\right)\varphi(r) = 0 \tag{11}$$

$$\left(\frac{d}{dr} + \frac{m+1}{r} - \frac{M}{\hbar v_F}\right)\chi(r) = 0 \tag{12}$$

There are two possibilities, depending on the orientation of the local magnetization: When the magnetization is oriented clockwise, $\vec{M} < 0$, the physical zero-modes read:

$$\psi(\vec{r}) = a_m \begin{pmatrix} 0 \\ 1 \end{pmatrix} \frac{e^{-\frac{|M|}{\hbar v_F} r}}{r^{m+1}} e^{i(m+1)\phi}\,, \tag{13}$$

where a_m is the normalization constant give by $a_m = \frac{1}{\sqrt{2\pi(|2m+1|!)}}\left(\frac{2M}{\hbar v_F}\right)^{|m+1/2|+1/2}$. Solution (13) can be normalized only if $2m+1 < 0$; in addition, these solutions are spin-polarized with $s_\phi = -\frac{\hbar}{2}$, so that the vortex provided a mechanism for supporting spin-polarized zero-energy current on the TI surface (see Fig. 2). On the other hand, if the local magnetization is oriented counterclockwise, $\vec{M} > 0$, the zero-mode solutions are spin polarized with $s_\phi = +\frac{\hbar}{2}$ and given by:

$$\psi(\vec{r}) = b_m \begin{pmatrix} 1 \\ 0 \end{pmatrix} r^m e^{-\frac{|M|}{\hbar v_F} r} e^{im\phi}\,, \tag{14}$$

with $b_m = \frac{1}{\sqrt{2\pi(2m+1!)}}\left(\frac{2M}{\hbar v_F}\right)^{m+1}$. This solution has physical meaning only if $2m+1 > 0$. Both sets of spin-polarized zero-modes carry integer electronic charge:

$$Q/e^- = \int d^2r\, j^0 = \int d^2r \psi^\dagger \psi = 1\,. \tag{15}$$

Physically, it is precisely the coupling between the orbital angular momentum of the carriers and the magnetic moments of the vortex that yields polarized charged zero-modes confined to a circular region with radius $r_0 \equiv \hbar v_F/|M|$, around the vortex center. Taking Sb_2Te_2 as an example, one has $\hbar v_F \approx 3.7\text{eV Å}$ while $|M|$ is typically $\sim 50\text{meV}$ giving us $r_0 \approx 74\text{Å}$ (see Fig.2). It is important to remark that, in this case, there is only one fermion per quantum state (which is denoted by the quantum number m); indeed, as we have seen, the vortex chirality requires that these states are polarized. Therefore, in principle, such zero modes may contain a very large number of polarized carriers (one per each possible m) for a given vortex chirality.



$M < 0, \quad S_\phi = -\hbar/2$ $\quad$ $M > 0, \quad S_\phi = +\hbar/2$

FIG. 2. (Color online) By breaking time reversal symmetry a ferromagnetic vortex forces the electrons spins, confined to the topological insulator surface, to align along its magnetization direction, according to Eqs. (13) and (14) . Such an alignment is fully accomplished by zero-mode fermions that appear concentrated in a region of radius r_0 around the vortex center, what yields the possibility of highly spin-polarized zero-mode current.

IV. BOUND STATES

In addition to the zero-modes at there also exist additional bound-state solutions of Eqs. (9) and (10) with $E < 0$. The solutions correspond to internal modes in which the electrons wave-functions undergo harmonically oscillations around the vortex center. Bound-states can be induced in 2D or 3D TI's by vacancies or holes in the band gap. These defects can affect the transport properties of TI's via additional bound-states near the system boundary due to the coexistence of the in-gap bound-states near the edge or surface states[47]. Here, this effect arises due to the exchange coupling between the spin of the surface carriers and the magnetic moment of the vortex in the surface.

To obtain the bound-states we proceed in the usual way by looking for solutions of Eqs. (9) and (10) with negative energy, $E < 0$. Decouple these equations we obtain the following second order differential equations for the up component of the spinor (down component satisfies a similar eq., as below):

$$\frac{d^2\varphi(r)}{dr^2}+\frac{1}{r}\frac{d\varphi(r)}{dr}+\left(-\frac{m^2}{r^2}+\frac{M}{\hbar v_F}\frac{2m+1}{r}+\frac{E^2-M^2}{\hbar^2 v_F^2}\right)\varphi(r)=0\,. \tag{16}$$

To solve this equation, we use the correspondence between this problem and the 2D quantum Kepler problem (2D hydrogen atom). Such an equation admits solutions for electrons restricted to move in a plane around the nucleus, due to the potential $1/r$ with[48] $r^2=x^2+y^2$. The energy eigenvalues for the bound-states read:

$$E^2(n\,,\,m)=M^2\left[1-\frac{(m+1/2)^2}{(n+|m|+1/2)^2}\right], \tag{17}$$

with $n=1\,,2\,,3\,,\ldots$ and $|m|=0\,,1\,,2\,,\ldots\,n-1$. The radial eigenfunction can be obtained in terms of the hypergeometric function[48]:

$$\varphi_{n,m}(r)=\frac{\alpha_{n,m}}{(2|m|)!}\left[\frac{(n+|m|-1)!}{(2n-1)(n-|m|-1)!}\right]^{1/2}\times (\alpha_{n,m}r)^{|m|}e^{-\alpha_n r/2}\times {}_1F_1(-n+|m|+1,\,2|m|+1,\,\alpha_{n,m}r)\,. \tag{18}$$

Here, $\alpha_{n,m}=\frac{M(2m+1)}{(n-1/2)\hbar v_F}$ and ${}_1F_1$ is a confluent hypergeometric function. The component $\chi_{n,m}(r)$ of the spinor can be obtained from the $\varphi_{n,m}(r)$ by using equation (9).



FIG. 3. (Color online) The 'atomic' energy levels allowed for zero-modes and bound-states in the composed vortex-topological insulator system. An almost continuum spectrum is allowed between $-|M|$ and $\approx -0.94|M|$, where the most energetic bound-state occurs, while $\delta\approx +0.94|M|$ gaps this state from zero-modes.

Note that the infinite number of energy levels goes from $-|M|$ (the ground state) to $-2\sqrt{2}|M|/3\approx -0.9428|M|$ (the most excited state) but, in contrast to the $2D$ Hydrogen atom, the ground state is characterized by $n=\infty$ while the most excited state is characterized by $n=1$ (Fig. 3 illustrate the energy levels for such an 'atom'). The Bohr radius is $a_0=\frac{2\hbar v_F}{|M|}$, which is on the order of $148\mathring{A}$ for typical values of v_F and $|M|$. The most probable radius for these bound-states are $(n+1/2)a_0$ (let us recall that a_0 is much smaller than the size of typical magnetic nanodisks containing a vortex ground state. This is particularly important if these nanoscaled systems are employed for TI coating). In addition, the most excited bound-state radial wavefunction are:

$$\varphi_{10}(r)=\frac{2M}{\hbar v_F}e^{-|M|r/\hbar v_F}\,, \tag{19}$$

$$\chi_{10}(r)=\pm\frac{3i}{\sqrt{2}}\left(\frac{2M}{\hbar v_F}+\frac{m}{r}\right)e^{-|M|r/\hbar v_F}\,, \tag{20}$$

from what becomes more evident that the effective potential brought about by the magnetic vortex is able to confine the TI surface electrons in a limited region around the vortex center. Similarly to the zero energy modes, the region of large probability is a circle with a radius on the order of $\hbar v_F/|M|$, depending only on the value of the local magnetization M, decreasing as $|M|$ gets higher. In addition, bound-states with normalized wavefunctions are possible only if $M(2m+1)>0$. Therefore, there are two possibilities: if $M>0$ then $m>-1/2$, while for $M<0$, $m<-1/2$ must be taken. As a consequence, all bound electrons move around the same sense, determined by the vortex magnetization chirality (see Fig. 2).

Besides the bound-states discussed in the former sections, a continuum spectrum of scattered states should takes place; they could be parametrized by a continuum wave-vector $\vec{k}$ whose associates wave-functions being solutions of equation (16). For these states, the relevant results can be obtained in terms of the phase-shifts of the carriers, cross-section and resistivity, and will be presented elsewhere[49].

Some additional remarks are now in order. In general, the Dirac equation has negative and positive energy solutions. The negative energy solutions correspond to the states in the valence band while the positive energy solutions correspond to the states in the conduction band (considering the chemical potential located at $E=0$). In the ground-state, all the negative energy levels are filled and, therefore, the ground-state is degenerate (since the zero modes energy are degenerate). Then, what is the ground-state at TI surface state in the presence of a magnetic vortex? In other words, how could one treat the zero-energy modes whenever constructing the ground-state of the TI-ferromagnetic composed material? Should the states be empty like the case of positive energy or should they be filled as they are for negative energy states?

If they are filled (or partially filled) when an electric current is established over the TI with a vortex (and only zero-modes are excited), then the current is spin polarized, being the polarization given by sense of the magnetization. However if the bound-states are excited to the conduction band (overcoming $\delta = 2\sqrt{2}|M|/3 \approx 0.9428|M|$), then the current has a gap and the current will display steps when increasing the external electrical field similar to the Hall effect under increasing of the external magnetic field.

If the previous study were performed for a planar radial vortex with magnetization exhibiting radial-like profile, $\phi_0 = 0$, we should have realized that the zero-modes were not spin polarized while all bound-states, with $E < 0$, were ruled out.

It is noteworthy to mention that the authors of Ref. [50] have treated a similar problem of a magnetic vortex coating the surface of a TI. However, in their work they have considered the question of whether an effective gauge field would be induced by strain on the surface of a topological insulator. They have shown that the gauge fields arise when the surface is coated with an easy-plane ferromagnet with magnetization parallel to the surface[50].

V. DISCUSSIONS AND CONCLUSIONS

In summary, we have investigated the effects of a closed flux magnetization (in-plane vortex with $\phi_0 = \pi/2$) on the surface of a $3D$ strong topological insulator with one Dirac cone. We have shown that the interplay between topologically protected surface states and the broken of time-reversal symmetry by an in-plane magnetic vortex coating TI surface does not generate a mass gap; It rather leads bound-states and localized spin polarized charged zero-modes. Indeed, considering the bound-states, the very presence of the vortex generates an effective potential for the TI surface carriers which is equivalent to the problem of a two-dimensional hydrogen atom. The range of energy levels of the bound-states is very short and goes from $-\mid M\mid$ to $-0.9428|M|$ (comprising infinite possible states in this range), which is energetically separated from the zero-modes, $E = 0$, by a large gap $\delta \approx 0.9428|M|$; a picture of such energetics may be seen in Fig. 3). Zero-modes and bound-states decay exponentially as $\exp(-|M|r/\hbar v_F)$ away from the vortex center, which means that some electrons are essentially localized in a circular region with radius $r_0 = \hbar v_F/|M|$ (around 70Å for typical TI's with "atomic" binding energy $\sim 50meV$. Physical systems supporting stable zero-modes deserve special attention, for instance, in topological quantum computation for the ground-state degeneracy remains unaffected by small perturbations, what can render error free quantum information encoding.

Instead of large magnetic films, nanoscale magnets can also be used to form the composed system with TI's. In this case, special care must be taken because vortices in confined structures must have out-of-plane components near its center, what opens up a gap in the TI electronic states. The remaining effects of a vortex, as studied here, should remain qualitatively unaffected, still supporting zero-modes and bound-states on the TI surface. Even in the case of nanomagnets, in-plane vortices may take place, for instance, if a cavity is intentionally inserted at the center of the disk[33,34] or we employ other geometries, such as a nanotorus[51]. Another interesting possibility is to coat TI surface with an elliptic thin disc supporting a pair vortex-antivortex, with opposite chiralities. On the TI surface it would be induced two distinct zero-modes fermionic current flowing in opposite directions, around each of the vortex center.

Finally, it should be remarked that the detection of the electronic modes predicted here on the surface of a magnetically coated TI could be an interesting alternative method to experimentally detect vortices in layered 2D Heisenberg ferromagnets.

Acknowledgments

The authors thank O.M. Del Cima, A.H. Gomes and D.H.T. Franco for useful discussions. They are also grateful to CNPq, FAPEMIG and CAPES (Brazilian agencies) for financial support.

---

[*] jakson.fonseca@ufv.br
[†] winder@ufv.br
URL: https://sites.google.com/site/wamouramelo/home
[‡] apereira@ufv.br
URL: https://sites.google.com/site/quantumafra/

[1] D. Hsieh, D. Qian, L. Wray, Y. Xia, Y. S. Hor, R. J. Cava, and M. Z. Hasan, Nature **452**, 970 (2008).

[2] Y.L. Chen, J.G. Analytis, J.-H. Chu, Z.K. Liu, S.-K. Mo, X.L. Qi, H.J. Zhang, D.H. Lu, X. Dai, Z. Fang, S.C. Zhang, I.R. Fisher, Z. Hussain, Z.-X. Shen , Science **325**, 178 (2009).

[3] Y. Xia, D. Qian, D. Hsieh, L. Wray, A. Pal, H. Lin, A. Bansil, D. Grauer, Y. Hor, R.J. Cava, M. Z. Hasan, Nature Phys. **5**, 398 (2009).

[4] S. Chadov, X.L. Qi, J. Kúbler, G.H. Fecher, C. Felser, S.C. Zhang, Nature Mater. **9**, 541 (2010).

[5] L.A. Wray, S.-Y. Xu, Y. Xia, Y.S. Hor, D. Qian, A.V. Fedorov, H. Lin, A. Bansil, R.J. Cava, M.Z. Hasan, Nature Phys. **6**, 855 (2010).

[6] A.H. Castro-Neto, F. Guinea, N.M. Peres, K.S. Novoselov and A.K. Geim, Rev. Mod. Phys. **81**, 109 (2009).

[7] For reviews, see X.L. Qi and S.C. Zhang, Phys. Today **63**,

33 (2010); M.Z. Hasan and C.L. Kane, Rev. Mod. Phys. **82**, 3045 (2010) and X.L. Qi, S.C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011)

[8] Y.L. Chen, J.-H. Chu, J. G. Analytis, Z.K. Liu, K. Igarashi, H.-H. Kuo, X.L. Qi, S.K. Mo, R.G. Moore, D.H. Lu, M. Hashimoto, T. Sasagawa, S.C. Zhang, I.R. Fisher, Z. Hussain, and Z. X. Shen, , Science **329**, 659 (2010).

[9] Q. Liu, C.-X. Liu, C. Xu, X.-L. Qi, and S.-C. Zhang, Phys. Rev. Lett. **102**, 156603 (2009).

[10] A.M. Essin, J. E. Moore, and D. Vanderbilt, Phys. Rev. Lett. **102**, 146805 (2009).

[11] X.-L. Qi, T. Hughes, and S.-C. Zhang, Phys. Rev. B **78**, 195424 (2008).

[12] I. Garate and M. Franz, Phys. Rev. Lett. **104**, 146802 (2010).

[13] K. Nomura and N. Nagaosa, Phys. Rev. B **82**, 161401 (2010).

[14] D.-X. Qu, et al., Science **329**, 821 (2010).

[15] L. A. W. et al., Nat. Phys. **7**, 32 (2010).

[16] Q. Liu, C.-L. Liu, C. Xu, X.-L. Qi, and S.-C. Zhang, Phys. Rev. Lett. **102**, 156603 (2009).

[17] A.R. Pereira and W.A. Moura-Melo, "*Vortex behavior in ferromagnetic systems with small defects: From macro to nanostructured magnets*", in "Electromagnetic, Magnetostatic, and Exchange-Interaction Vortices in Confined Magnetic Structures", Edited by E.O. Kamenetskii, Transworld Research Network, Kerala, India (2008).

[18] G.M. Wysin, Phys. Rev. B **94**, 8780 (1994).

[19] F.G. Mertens, A.R. Bishop, G.M. Wysin, and C. Kawabata, Phys. Rev. B **39**, 591 (1989).

[20] A.S.T. Pires, A.R. Pereira, and M.E. Gouvêa, Phys. Rev. B **49**, 9663 (1994).

[21] M.T. Hutchings, P. Day, E. Janke, and R. Pynn, J. Magn. Magn. Mater. **54-57**, 673 (1986).

[22] C.E. Zaspel, T.E. Grigereit, and J.E. Drumheller, Phys. Rev. Lett. **74**,4539 (1995).

[23] A.R. Pereira and J.E.R. Costa, J. Magn. Magn. Mater. **162**, 219 (1996).

[24] D.A. Dimitrov and G.M. Wysin, Phys. Rev. B **53**, 8539 (1996).

[25] M.M. Bogdan and C.E. Zaspel, Phys. St. Solidi. A **189**, 983 (2002).

[26] A.R. Pereira, S.A. Leonel, P.Z. Coura, and B.V. Costa, Phys. Rev. B **71**, 014403 (2005).

[27] F.M. Paula, A.R. Pereira, and L.A.S. Mól, Phys. Lett. A **329**, 155 (2004).

[28] F.M. Paula, A.R. Pereira, and G.M. Wysin, Phys. Rev. B **72**, 094425 (2005).

[29] A.R. Pereira and G.M. Wysin, Phys. Rev. B **73**, 214402 (2006).

[30] A.R. Pereira, L.A.S. Mól, S.A. Leonel, P.Z. Coura, and B.V. Costa,Phys. Rev. B **68**,132409 (2003).

[31] J. Miltat and A. Thiaville, Science **298**, 555 (2002).

[32] K.Y. Guslienko, X.F. Han, XF, D.J. Keavney, R. Divan, and S.D. Bader, Phys. Rev. Lett. **96**, 067205 (2006).

[33] M. Rahm, J. Biberger, V. Umansky, and, D. Weiss, J. Appl. Phys. **93**, 7429 (2003).

[34] M. Rahm, R. Hollinger, V. Umansky, and, D. Weiss, J. Appl. Phys.**95**, 6708 (2004).

[35] A.R. Pereira, Phys. Rev. B **71**, 224404 (2005).

[36] A.R. Pereira, J. Appl. Phys. **97**, 094303 (2005).

[37] R.L. Silva, A.R. Pereira, R.C. Silva, W.A. Moura-Melo, N.M. Oliveira-Neto, S.A. Leonel, and P.Z. Coura, Phys. Rev. B **78**, 054423 (2008).

[38] S.-Y. Xu, et.al., Nature Phys. **8**, 616 (2012).

[39] A. Zazunov, A. Kundu, A. Hütem and R. Egger, Phys. Rev. B **82**, 155431 (2010).

[40] D. A. Abanin and D. A. Pesin, Phys. Rev. Lett. **106**, 136802 (2011).

[41] T.M. Schmidt, R.H. Miwa and A. Fazzio, Phys. Rev. B **84**, 245418 (2011).

[42] J.M. Fonseca, W.A. Moura-Melo and A.R. Pereira, J. Appl. Phys. **111**, 064913 (2012).

[43] R. Jackiw, *"Fractional and Majorana Fermions: The Physics of Zero Energy Modes"*, arXiv: 1104.4486v1.

[44] H. Zhang, C.-X. Liu, X.-L. Qi, X. Dai, Z. Fang, and S.-C. Zhang, Nature Phys. **5**, 438 (2009).

[45] J.K. Pachvos and M. Stone, *"The index theorem for graphene"*, arXi:cond-mat/0607394v4 (2007).

[46] J.M. Fonseca, *"Topologically protected surface states and index theorem for topological insulators"* work in progress.

[47] W.Y. Shan, J.L., H.Z. Lu and S.Q. Shen Phys. rev. B **84**, 035307 (2011).

[48] X.L. Yang, S.H. Guo, F.T. Chan, K.W. Wong and W.Y. Ching, Phys. Rev. A **43**, 1186 (1991).

[49] J.M. Fonseca, W.A. Moura-Melo and A.R. Pereira, work in progress.

[50] M.I. Katsnelson, F. Guinea and M.A.H. Vozmediano, *"Gauge fields at the surface of topological insulators"*, arXix:cond-mat/1105.6132v2 (2011).

[51] V.L. Carvalho-Santos, W.A. Moura-Melo, and A.R. Pereira, J. Appl. Phys. **108**, 094310 (2010).